

Review paper

Cybersickness: A multisensory integration perspective

Maria Gallagher and Elisa Raffaella Ferrè

Department of Psychology, Royal Holloway University of London, Egham, UK

Corresponding Author:

Elisa Raffaella Ferrè

Department of Psychology

Royal Holloway University of London

Egham, Surrey TW20 0EX, UK

Telephone: +44 (0) 1784 443530

Email: E.Ferre@rhul.ac.uk

Summary

In the past decade, there has been a rapid advance in Virtual Reality (VR) technology. Key to the user's VR experience are multimodal interactions involving all senses. The human brain **must** integrate real-time vision, hearing, vestibular and proprioceptive inputs to produce the compelling and captivating feeling of immersion in a VR environment. A serious problem with VR is that users **may** develop symptoms similar to motion sickness, a malady called *cybersickness*. **At present the underlying cause of cybersickness is not yet fully understood.** **Cybersickness may be due to a discrepancy between the sensory signals which provide information about the body's orientation and motion: in many VR applications, optic flow elicits an illusory sensation of motion which tells users that they are moving in a certain direction with certain acceleration. However, since users are not actually moving, their proprioceptive and vestibular organs provide no cues of self-motion. These conflicting signals may lead to sensory discrepancies and eventually cybersickness.** Here we review the current literature to develop a **conceptual scheme** for understanding the neural mechanisms of cybersickness. **We discuss an approach to cybersickness based on sensory cue integration, focusing on the dynamic re-weighting of visual and vestibular signals for self-motion.**

Keywords

Cybersickness, Virtual Reality, Motion Sickness, Vestibular System, Multisensory Integration

Introduction

Virtual Reality (VR) came to the public's attention in the late 1980s. However, the real growth in popularity of VR has been observed in more recent years, when the development in technology allowed a user to be immersed in a completely virtual world through the use of 3D real-time computer graphics and advanced display devices. A report by Goldman Sachs in January 2016 predicted that revenue from VR would approach \$80 billion by 2025, with 315 million users (Bellini et al. 2016). Pivotal to the user's VR experience are multimodal interactions involving all senses. The human brain must integrate real-time vision, hearing, vestibular, and proprioceptive inputs to produce a compelling and captivating feeling of immersion and presence in a VR environment akin to real life scenarios (Azmandian et al. 2016; Herbelin et al. 2015). VR has proved to be beneficial in several applications ranging from educational and training platforms, to recreational gaming or media viewing, as well as flight simulators and medical rehabilitation (Alaker et al. 2016; Pelargos et al. 2016; Valmaggia et al. 2016; Viñas-Diz and Sobrido-Prieto 2016).

A troublesome problem with VR is that between 20% and 80% of users exhibit symptoms that parallel symptoms of classical motion sickness (Cobb et al. 1999; Munafo et al. 2017). This so-called *cybersickness* can be profoundly unsettling, and may compromise well-being and performance in VR training (Cobb et al. 1999; Sharples et al. 2008; Fiore et al. 2013; Llorach et al. 2014; Munafo et al. 2017). Symptoms of cybersickness include discomfort, apathy, nausea, drowsiness, disorientation, eyestrain and fatigue (Stanney et al. 1997; Rebenitsch and Owen 2016). Although its causes are not yet entirely clear, cybersickness occurrence might be dependent on different factors, including the user's gender (Munafo et al. 2017), age (Arns and Cerney 2005), duration of VR exposure (Liu 2014), and hardware issues,

1 such as lag (i.e., delays in updating the visual scene following a user's movement or action) or
2 flicker (i.e., flashing of the visual scene relating to refresh rates) (Moss et al. 2011).
3

4
5 In the past decade, there has been a rapid advance in VR to increase the simulation
6 realism by means of integrated multimodal experiences (i.e. 3D audio, haptic and force
7 feedback, etc.). Significant technical improvements have also occurred, including better
8
9 positional tracking, display resolution, and display refresh rate (Harsora et al. 2017). For
10
11 example, many Head Mounted Displays (HMDs) (i.e., Oculus Rift CV1 and HTC Vive) are
12
13 capable of tracking the position of the head in space via numerous sensors and cameras, have a
14
15 refresh rate of 90Hz, and visual resolution of 1080x1200 resolution for each eye. The
16
17 improvement in VR technology is therefore self-evident, and even more if compared to earlier
18
19 versions of commercial HMDs (i.e., no positional tracking capabilities, a 60Hz refresh rate,
20
21 and resolution of 640x800 for each eye) (Harsora et al. 2017). Moreover, the earliest HMDs
22
23 had much more limited technical capabilities: 30 frames per second, a 40 degree field of
24
25 view, and rather cumbersome head position sensors (Sutherland 1968). Further
26
27 improvements are likely in the coming years, such as foveated rendering and adaptive displays
28
29 which increase the resolution and refresh rate of the display at selected locations of the VR
30
31 scene (Padmanaban et al. 2017; Lee et al. 2017; Hsu et al. 2017), increasing realism and
32
33 immersion in VR (see Bastug et al. 2017 for a review). Despite these advances, there are still
34
35 some issues in VR and as one leading iconoclast in VR technology - Palmer Luckey - has
36
37 indicated, "*VR isn't perfect right now*" (Kushner 2016). Cybersickness remains a barrier to VR
38
39 use: it has even been argued that the more realistic the VR environment, the more likely it is
40
41 that the user experiences cybersickness (Merhi et al. 2007; Davis et al. 2015). Although the
42
43 exact reason behind this is still unclear, better resolution of optic flow might potentially
44
45 exacerbate visual-vestibular-proprioceptive conflicts (Kennedy et al., 2003; Merhi et al.,
46
47 2007; Davis, Nesbitt and Nalivaiko, 2015). Alternatively, one can hypothesise that the
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 increased feeling of presence (i.e. the feeling of “being there” in a VR environment) due to
2 the availability of detailed simulated environments might exacerbate cybersickness
3 symptoms (Kennedy et al., 2003; Merhi et al. 2007; Davis et al. 2015). In such cases, it is
4 possible that the detailed, realistic environment and increased presence may render the
5 conflict in sensory signals more dramatic, increasing cybersickness. Further research is
6 necessary to explore the exact relationship between increased feeling of presence and
7 cybersickness.

16 The root causes of cybersickness remain poorly understood, however more general
17 accounts of motion sickness have been adapted as explanations given the similarity between
18 cybersickness and other types of motion sickness. These theories include Reason and Brand's
19 (1975) Neural Mismatch Theory, based on sensory conflict, and Riccio and Stoffregen's (1991)
20 Postural Instability Theory based on movement control. However, neither theory can fully
21 explain why cybersickness arises, nor can they predict an individual's likelihood of developing
22 symptoms. Despite these theories highlighting the role of multiple sensory inputs in the cause
23 of cybersickness, more recent advances in knowledge of multisensory integration have only
24 recently applied to this field (Balter et al. 2004; Oman 2012; Jürgens et al. 2016; Weech and
25 Troje 2017).

41 Here we review the current literature to provide an overview of cybersickness and
42 perceptual after-effects induced by VR, as well as evaluating the current theories of
43 cybersickness. We highlight how dynamic re-weighting of sensory cues can be implicated in
44 cybersickness development and after-effects of VR exposure. This approach may provide
45 insight into cybersickness, facilitating further research and leading to developments which
46 may enhance the VR experience.

The vestibular system and sensory integration for self-motion perception

In everyday life, our perception of self-motion depends on the coherent integration of visual, proprioceptive and vestibular information. When we move in the surrounding environment, the visual system provides retinal-image motion (optic flow) cues, the proprioceptive receptors in the muscles, tendons, and joints sense the relative position of body parts in space and the vestibular system encodes angular and linear acceleration.

Since Gibson in 1950, motion related retinal-images have been considered essential for eliciting sensations of self-motion displacement (Gibson, 1950). However, it is now recognised that extra-visual cues make an equal contribution to self-motion perception. In the absence of visual cues, humans rely on vestibular information to estimate bodily motion (Israël and Berthoz 1989; Irsaël et al. 1993; Berthoz et al. 1995). The vestibular system is a set of sensory organs located in the inner ear, comprising of three orthogonal semicircular canals (anterior, posterior and horizontal) that sense rotational acceleration of the head in three-dimensional space and around three cardinal axes (yaw, roll, pitch), and two otolith organs (utricle and saccule) that code translational acceleration, including the orientation of the head relative to gravity. Vestibular organs are extremely sensitive to even the slightest changes in rotation and linear movement of the head, providing the feedback necessary for the brain to dictate adjustments that allow the body to maintain balance. Dynamic vestibular inputs from the semicircular canals are associated with low-level visuo-vestibular interactions to control gaze and eye fixation, while gravitational inputs from the otolith organs contribute to path integration and navigation. Thus, vestibular cues are fundamental for perception of self-motion, aiding us in distinguishing self from object motion and providing us with a sense of where we are in space (Green and Angelaki 2010; Greenlee et al. 2016). Thus, it might not be surprising that the vestibular system plays a vital contribution in the development of motion

sickness, including cybersickness. Accordingly, labyrinthine-defective patients do not experience any motion sickness symptoms (Cheung et al. 1991; Paillard et al. 2013) while blind individuals do, suggesting that visual information, while implicated motion sickness, is not as crucial as vestibular signalling (Greybiel 1970).

Multimodal interactions between visual, somatosensory, proprioceptive and vestibular signals have been described in almost all vestibular relays, including the vestibular nuclei, the thalamus and several areas in the cerebral cortex (Lopez et al. 2012; Zu Eulenburg et al. 2012). Electrophysiological studies have identified a widespread vestibular network in which the core area is the Parieto Insular Vestibular Cortex (PIVC) (Guldin and Grüsser 1998; Chen et al. 2010). This area consists of the posterior insula/retroinsular cortex in the bank of the lateral sulcus (Guldin and Grüsser 1998). The human homologue of the primate PIVC is a distributed set of regions, including retroinsular cortex, temporoparietal junction and somatosensory cortices (Fasold et al. 2002; Lopez and Blanke 2011; Zu Eulenburg et al. 2012). Neuroimaging studies with artificial vestibular stimulation showed activation in large swathes of the cortex, including activations of classically unimodal sensory and motor regions, supporting function integration between signals and inputs from other senses (Ferrè and Haggard 2015).

Multisensory interactions between sensory modalities are fundamental in shaping our perceptual experiences. Sensory signals presented simultaneously in more than one sensory channel tend to be detected more accurately than the same signals presented individually (Stein et al. 1996). Most studies of vestibular-multisensory interactions have focused on *multisensory convergence* between vestibular and visual signals. Multisensory convergence involves multisensory integration of different signals related to a common external source object or percept (Ernst and Bühlhoff 2004). Critically, the neural mechanism underlying multisensory convergence is likely to be a process seeking to reduce perceptual uncertainty about the source

(Knill and Pouget 2004), which often involves optimal combination of cues across modalities (Ernst and Banks 2002).

The predominant theme in recent electrophysiological work has been the convergence between vestibular signals and visual signals for perception of self-motion, spatial orientation, and navigation in the environment. Multisensory neurons coding for visual and vestibular signals have been described in the macaque ventral intraparietal area (VIP, Bremmer et al. 2002), which is considered homologous to human vestibular areas in the posterior parietal cortex (Lopez and Blanke 2011). Visuo-vestibular interactions are often interpreted within the framework of optimal cue combination for multisensory perception of a single underlying quality (Gu et al. 2008; Fetsch et al. 2009). For instance, heading direction is accurately perceived in visuo-vestibular multimodal conditions. Macaques trained to complete a heading discrimination task where cues were provided by an optic flow (vision), or by a motion platform (vestibular), or combined (visuo-vestibular), showed smaller thresholds for detecting head direction under the combined sensory condition than either of the unimodal visual or vestibular conditions (Gu et al. 2008). Importantly, these results were mirrored by the activity of dorsal medial superior temporal (MSTd) neurons: the neurons' preferred heading direction was similar for both visual and vestibular modalities, suggesting a neural mechanism for perceptual integration. Fetsch et al. (2009) further investigated this integration by dynamically modifying the reliability of the visual cue by reducing optic flow coherence and placing the visual and vestibular cues in conflict. The results matched the predictions of optimal cue integration; as the coherence of the visual cue decreased and became less reliable, the weighting of the vestibular cue increased. Interestingly, the conflict between the visual and vestibular cues did not prevent sensory integration. Similar results were found in human participants. Thus, visual and vestibular signals might be combined optimally when estimating heading direction – and therefore self-motion (however see de Winkel et al. 2010; de Winkel et al. 2013

for contrary findings). The activity of neurons in MSTd is a likely neural mechanism for this effect (Angelaki et al. 2011).

What is cybersickness?

Cybersickness is an unpleasant sensation, comprising of symptoms of disorientation, drowsiness, eyestrain, and nausea arising from exposure to immersive VR environments. Cybersickness is triggered by visually-induced illusory motion within an immersive VR environment, in which an optic flow provides motion information in the absence of corresponding vestibular signals (Reason and Brand 1975; Hill and Howarth, 2000; Keshavarz et al. 2015; Rebenitsch and Owen 2016). For this reason, cybersickness is slightly different compared to traditional motion sickness syndromes, such as car-sickness, sea-sickness and air-sickness, in which the physical movement of the vehicle triggers motion sickness symptoms (Reason and Brand, 1975; Golding 2016). However, given its similarity to motion sickness, cybersickness can be regarded as a type of visually-induced motion sickness.

Many VR applications employ an optic flow pattern which elicits an illusory feeling of self-motion, namely *vection*¹. In a VR driving simulator, for instance, the simulation provides accurate optic flow patterns of the road, buildings and other parts of the environment, eliciting clear vection sensations. The visual signals tell the user that they are moving in a certain direction with a certain acceleration. However, since the user is not actually moving, the vestibular organs provide no cues for linear or angular acceleration. As visual signals for self-motion are not corroborated by inertial forces transmitted through the vestibular system, a

¹ Although vection is a complex perceptual experience (see Palmisano et al. 2015 for a review), for the purposes of this review vection is defined as a sensation of self-motion arising from visual cues in a VR environment.

1 sensory conflict is likely to occur, and subsequently lead to cybersickness (Keshavarz et al.
2 2015).
3

4 The example above refers to the scenario in which the user does not move their head
5 during VR exposure. However, many VR HMDs are now supplied with positional trackers,
6 which enable the user to physically move in the real world while exploring the VR environment
7 (Harsora et al. 2017). When physically moving in VR, visual cues are supported by vestibular
8 information. This may drastically reduce the conflict between sensory modalities, and may
9 prevent the occurrence of cybersickness. At present, few empirical studies have compared
10 levels of cybersickness between locomotion techniques, and findings are somewhat mixed
11 (Chance et al. 1998; Zambaka et al. 2004; Peck et al. 2011; Llorach et al. 2014). For example,
12 Llorach et al. (2014) compared navigation in a VR environment via a game controller, in which
13 the user remained stationary and moved via controlling joysticks, and a position estimation
14 system which tracked the users' actual movements. Levels of cybersickness were much higher
15 when the user did not move and used the game controller compared to when they navigated by
16 physically moving. By contrast, Peck et al. (2011) found no differences in cybersickness
17 symptoms between VR scenarios in which users freely moved, walked in place, or explored the
18 VR environment using a joystick, despite finding better performance in the free movement
19 scenario. Although new developments in positional trackers also allow users to explore the VR
20 environment through head movements while sitting,vection appears much stronger when users
21 move their heads, and possibly contributing to cybersickness (Ash et al. 2011). In addition, it is
22 not possible to exclude that incorrect updating of the visual scene during active head motion
23 might elicit cybersickness (Ash and Palmisano 2012). For example, cybersickness increases
24 when the visual VR scene moves in the same direction of the physical head movement
25 (Palmisano et al. 2017). Similar conflicts between visual and vestibular signals are implicated
26 in car-, sea-, and space-sickness (Lackner 2014).
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Traditionally, the presence of cybersickness has been explored via subjective self-reports. The Simulator Sickness Questionnaire (SSQ; Kennedy et al. 1993) is the most frequently used. This questionnaire breaks down motion sickness symptoms into three main categories: *disorientation* (D), including symptoms such as dizziness, vertigo and difficulty focusing, *oculomotor* (O), including eyestrain, headache, and blurred vision, and *nausea* (N), including stomach awareness, increased salivation, as well as nausea itself (Table 1). Cybersickness is characterised by severe and frequent disorientation symptoms, followed by nausea symptoms, and least oculomotor symptoms (a so-called D>N>O profile; Stanney et al. 1997; Rebenitsch and Owen 2016). This symptom profile further distinguishes cybersickness from other types of motion sickness. For example, simulator sickness has an O>N>D profile, sea-sickness an N>O>D profile, and space sickness an N>D>O profile (Stanney et al. 1997; Rebenitsch and Owen, 2016). Moreover, symptoms of cybersickness are reportedly much more severe than simulator sickness and other motion sickness symptoms (Kennedy et al. 2003).

*** PLEASE INSERT TABLE 1 HERE ***

Along with self-reported symptoms of cybersickness, a range of physiological changes have also been described, including increases in tachygastic power, heart rate and eyeblinks, decreases in bradygastic power, and changes in skin temperature and EEG power bands (Kim et al. 2005; Nalivaiko et al. 2015) (Table 1). Kim et al. (2001) identified increases in eyeblinks, skin conductance response, and heart rate, as well as decreases in the amplitude of photoplethysmogram associated with cybersickness. In addition, a selective modulation of the EEG gamma band activity was noted (Kim et al. 2001). Critically, physiological changes can predict cybersickness severity. For instance, bradygastic power, breathing rate, pulse amplitude and blinking rate can predict total or subscale scores of the SSQ (Dennison et al.

2016). Although physiological changes could represent an objective measure of cybersickness, they are not widely employed at present. Exploration of these changes may prove useful in determining the likelihood of cybersickness severity in individual users, therefore allowing intervention before cybersickness develops. An overview of cybersickness symptoms and physiological changes can be seen in Table 1.

The severity of cybersickness symptoms seems to be proportional to the duration of VR exposure: increasing exposure time within a single VR session is likely to increase cybersickness symptoms (Stanney et al. 2002; Moss et al. 2011). Liu (2014) found that participants completing a VR task had more severe symptoms as the duration increased from 5 to 15 minutes. Interestingly, a 20-minute session produced less severe symptoms than the 15-minute session, which might be due to adaptation to the VR environment (see below).

It is also possible that cybersickness symptoms do not recede immediately after cessation of VR, but rather linger on for some time following exposure. Stanney and Kennedy (1998) found cybersickness symptoms present at least one hour post-exposure. In accordance with the general profile of cybersickness, disorientation symptoms were most severe, followed by nausea and oculomotor symptoms. Startlingly, disorientation symptoms were almost 150 times higher immediately following VR exposure, and remained 95 times higher than the pre-exposure level one hour following exposure.

Although the profile of symptoms in cybersickness is relatively consistent, several factors influence whether an individual will develop the syndrome during VR exposure. Not surprisingly, people who have a history of motion sickness susceptibility are more likely to experience cybersickness, and are less likely to enjoy using the technology (Nichols 2000; Rebenitsch and Owen 2014). Age and gender also seem to be important: Arns and Cerney (2005) found that symptom severity and incidence increased with age, and a number of studies report that females are more susceptible to cybersickness than males (Stanney et al. 1999;

Stanney et al. 2003; Flanagan et al. 2005; Kim et al. 2008; Chen et al. 2015). The reason for this gender difference is somewhat unclear, and might be driven by the influence of the menstrual cycle (Clemes and Howarth 2005; however, Golding et al. 2005 argue that the influence of the menstrual cycle is too small to fully account for the gender difference), differences in postural stability between men and women (Koslucher et al. 2016), or a larger field of view in women (LaViola 2000).

Current interpretations of cybersickness

While an understanding of cybersickness and its influences is growing, uncovering precise neural mechanisms behind cybersickness is less straightforward. A specific framework for cybersickness has not been widely employed at present. Instead, general theories of motion sickness have been adopted to explain cybersickness (Table 2).

*** PLEASE INSERT TABLE 2 HERE ***

Reason and Brand (1975) suggested that a discrepancy between sensory modalities is the root cause of motion sickness syndromes. Two main forms of sensory conflict have been identified in motion sickness: intersensory conflicts between visual and vestibular signals, and intrasensory conflicts between the semicircular canals and otoliths within the vestibular system (Reason 1978). Thus, the vestibular system seems to be critical in causing motion sickness, as supported by evidence in peripheral vestibular patients who do not experience any form of motion sickness (Cheung et al. 1991; Paillard et al. 2013).

According to Reason and Brand's (1975) Neural Mismatch Theory, a copy of a self-generated movement is paired with the resulting sensory inputs to form a predicted pattern of

1 sensory cues, i.e. an engram (Reason, 1978). Then, a comparator module matches the actual
2 sensory inputs with the stored engrams. If the input and engram do not match, a discrepancy
3 arises and a mismatch signal is generated, triggering motion sickness. The strength of this
4 mismatch signal is dependent on how many sensory modalities are in conflict, the extent of the
5 discrepancy and the amount of previous exposure to the conflicting stimuli. Accordingly, the
6 strength of the mismatch signal corresponds to the latency and severity of motion sickness
7 symptoms. When VR users are immersed in applications where they perceive self-motion
8 through vection, visual signals suggesting movement conflict with vestibular inputs signalling
9 the user is stationary. A mismatch signal is then generated if no matching engram is found,
10 triggering cybersickness (Reason and Brand 1975; Reason 1978).
11
12
13
14
15
16
17
18
19
20
21
22
23

24 Reason (1978) proposed a further classification of sensory conflicts which have the
25 potential to trigger cybersickness. First, information signalled by visual and vestibular systems
26 is contradictory. For example, this may be the case when HMDs are improperly calibrated,
27 showing VR movements (visual cues) that are not properly aligned with the user's head
28 movements (vestibular cues). Second, visual information is not corroborated by expected
29 signals from the vestibular organs. This sensory conflict is unsurprisingly common in
30 cybersickness, when vection is not supported by vestibular information. Finally, vestibular
31 information is not corroborated by visual signals, as experienced in the use of HMDs without
32 head-tracking in which changes in head position may not be verified by changes in the VR
33 scene.
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 Although Reason and Brand's (1975) theory is widely accepted, it cannot fully account
49 for motion sickness, and therefore cybersickness, onset and development. First, this framework
50 lacks a clear physiological basis which would explain the importance of mismatch signals in
51 facilitating sickness (Oman 1988). Second, Reason and Brand's (1975) theory cannot account
52 for individual differences in motion sickness (Warwick-Evans et al. 1995; Davis et al. 2014).
53
54
55
56
57
58
59
60
61
62
63
64
65

For example, it is unclear why females should be more susceptible to motion sickness than males (Stanney et al. 1999; Stanney et al. 2003; Flanagan et al. 2005; Kim et al. 2008; Chen et al. 2015). Finally, the theory is unable to explain why some sensory cues are more likely to cause sickness than others. According to Neural Mismatch Theory, any sensory conflict triggering a mismatch signal should cause sickness, so it is unclear why particular stimuli are more nauseogenic than others. For example, scene oscillations within a VR environment are more likely to cause cybersickness than scenes with no oscillation (So and Lo, 1999; Lo and So, 2001). In particular, oscillations of around 0.2Hz in real motion sickness are highly nauseogenic, with oscillation along the fore-aft axis the most likely to cause sickness (Kennedy et al. 2010).

To address some of these open questions, Oman (1988) proposed that a desired body state prompts muscle activity and postural changes to reach that state. These changes provide signals which, along with external noise, are detected by different sensory modalities. An internal model based on all sensory modalities is formed, which is compared with actual sensory signals, providing a difference vector. Accordingly, greater sensory conflicts lead to a larger vector, which may reflect severe sickness.

Building upon the theory of Oman (1988), Bles et al. (1998) and Bos et al. (2008) proposed a more nuanced description of sensory conflict based on perception of the subjective vertical. The subjective vertical is formed from integrated sensory information from vision, proprioception, and the vestibular organs and is necessary for successful interactions with the external world (Barra et al. 2010). The visual and vestibular senses construct a model of the expected subjective vertical, as well as sensing the actual subjective vertical. The comparison between the sensed and expected verticals leads to a difference vector, prompting motion sickness. Motion sickness may therefore arise when there is an unexpected change in the subjective vertical, causing a conflict between the sensed and expected verticals. In the case of

cybersickness, the VR environment may contain aspects where the visual and vestibular vertical are at odds, however further research is necessary to unpack aspects of this theory of motion sickness.

While the underlying basis for sensory conflict-based models remains debated, empirical evidence for the involvement of sensory conflicts in motion sickness induced by physical movement (i.e., sea-sickness, car-sickness) highlights their contribution to the development of sickness symptoms (Kato and Kitazaki 2008; Wang and Lewis 2016; Wada et al. 2016; Wada and Yoshida 2016). The specific evidence for visuo-vestibular conflicts in cybersickness is growing, suggesting that the strength of the sensory conflict between visual and vestibular cues can lead to increased sickness (Bonato et al. 2009; Nishiike et al. 2013). For example, Akiduki et al. (2003) induced a visuo-vestibular conflict while participants were immersed in VR. Participants were instructed to follow a virtual ball around the room, allowing for a range of movements. The visuo-vestibular conflict was induced by doubling the range of movement of the VR environment background relative to the participants' head movements. Cybersickness symptoms were significantly greater during and immediately after VR exposure. In addition, more complex patterns of visual motion are also related to cybersickness. Keshavarz and Hecht (2011) found that rotation across two or three axes induced increased levels of sickness than a single axis of rotation. Thus, greater mismatches between visual and vestibular modalities might trigger symptoms of cybersickness, as purported by Reason and Brand (1975) and Oman (1988).

According to the above theories, sensory conflicts are the cause of motion sickness, however it is unclear how and why such conflicts would cause symptoms such as nausea. One well-cited hypothesis is that of Poison Theory (Treisman 1977). According to this hypothesis, sensory mismatches are part of an *early warning system* when an animal has ingested toxins. Nausea is therefore an adaptive, evolved response to sensory conflict aimed at ridding the

1 animal of dangerous toxins (Treisman 1977). While this hypothesis is a plausible explanation
2 for nausea symptoms of motion sickness, it does not account for other symptoms, such as
3 oculomotor or disorientation symptoms, and many authors argue that it is not a compelling
4 explanation for motion sickness (LaViola 2000; Oman 2012; Davis et al. 2015).
5
6
7

8
9
10 Riccio and Stoffregen (1991) proposed that motion sickness is a result of prolonged
11 postural instability: people are likely to suffer motion sickness when experiencing novel
12 situations for which they have not yet learned strategies to stabilise their posture (Stoffregen et
13 al. 2000; Villard et al. 2008). For example, Stoffregen and Smart (1998) found increases in
14 postural sway preceding symptoms of visually induced motion sickness when participants were
15 exposed to low-amplitude optical flow in an immersive environment. Moreover, Smart et al.
16 (2002) found that pitch velocity and vertical variability could predict which participants would
17 become sick when exposed to optic flow stimulation. However, the causal relation between
18 postural instability and cybersickness is not yet clear. For instance, Dennison and D’Zmura
19 (2017) found that postural sway was similar both before and during VR exposure, and
20 cybersickness increased both when participants were seated (and therefore unlikely to have an
21 unstable posture) and when they were standing (and thus subject to greater postural demands
22 and the potential for instability). Similarly, Warwick-Evans et al. (1995) found that motion
23 sickness was equally present when participants viewed a video while standing and while
24 restrained in a chair. Finally, Akiduki et al. (2003) demonstrated that postural instability (in
25 particular, body sway) was only significantly different post-exposure to VR, pointing to
26 instability as a consequence, rather than a cause, of cybersickness.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52

53 **Is it possible to prevent cybersickness?**
54
55
56
57
58
59
60
61
62
63
64
65

1 Techniques for preventing cybersickness include adaptation to VR through repeated
2 exposure (Barrett 2004; Keshavarz 2013), designing VR environments to include stable visual
3 references of the horizon or perceptual vertical (Han et al. 2011), developing applications based
4 on physical locomotion (Llorach et al. 2014), and providing concurrent vestibular signals by
5 means of galvanic vestibular stimulation (Cevette et al. 2012).
6
7
8
9
10

11 Adaptation to VR is arguably a more readily available technique to prevent
12 cybersickness, as it requires the user to repeatedly engage with VR, rather than modification of
13 VR applications or use of sensory substitution equipment (Keshavarz 2013; Golding 2016).
14
15
16
17
18

19 Several studies have shown reduced cybersickness following adaptation to VR: participants
20 exposed several times to the same VR scenario showed decreased cybersickness symptoms
21 (Regan 1995). Similarly, Hill and Howarth (2000) asked participants to complete five sessions
22 across five days in which they played a racing game for 20 minutes via a HMD. Some
23 participants also passively viewed the scene while the experimenter played the game, increasing
24 their exposure to VR. Participants experienced at least a mild degree of malaise during the first
25 session, however by the end of the fifth session seven out of 11 participants who played and
26 watched the game reported no cybersickness symptoms, suggesting faster adaptation to VR.
27
28
29
30
31
32
33
34
35
36
37
38

39 These findings are in accordance with the mentioned theories of motion sickness. For example,
40 Reason and Brand's (1975) Neural Mismatch theory predicts that as participants are further
41 exposed to conflicting sensory stimuli, their neural store creates a new engram, and in
42 subsequent exposures a mismatch signal is not generated to trigger cybersickness. Similarly, an
43 internal model of expected sensory signals within the VR environment may be updated as
44 participants are further exposed to the conflicting stimuli, as predicted by Oman (1988) and
45 Bles et al. (1998). By contrast, Riccio and Stoffregen's (1991) postural instability hypothesis
46 predicts that individuals adapt to motion sickness stimuli when they learn new strategies to
47 control their posture in the provocative environment. While adaptation to a cybersickness-
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 inducing environment is possible, it is yet unclear which of these motion sickness theories best
2 describes the process by which adaptation might occur.
3

4 Although the beneficial effects of VR adaptation on cybersickness symptoms are
5 promising (Howarth and Hodder 2008; Moss et al. 2011), several drawbacks are apparent.
6 First, repeated exposures are necessary for adaptation to be effective, which implies significant
7 commitment from VR users. Second, it is not yet clear how durable the benefits of adaptation
8 may be, limiting its utility. Finally, an inverse relation exists between reduction of
9 cybersickness through adaptation and the development of VR after-effects. These include
10 altered visual perceptions and balance problems (Stanney and Kennedy 1998; Stanney et al.
11 1999; Di Girolamo and Pic 2001; Harm et al. 2008). Thus, the more habituated an individual
12 becomes to the virtual world, the more likely they are maladapted to the real world on VR
13 cessation, incurring a range of after-effects (Wright 2014).
14
15

16 Sensory conflict appears the most likely explanation of cybersickness. Thus, a
17 reduction of these conflicts might prevent cybersickness. In VR the conflicting information
18 between vestibular and visual cues prevents the user from being able to accurately assess self-
19 motion. The presentation of 'rest frames' has been proposed as a method for reducing
20 cybersickness (LaViola 2000; Han et al. 2011). Rest frames are an explicit frame of reference
21 for spatial information concerning stationary objects, providing information on which to base
22 self-motion priors. For instance, cybersickness scores were reduced when users were exposed
23 to a VR rollercoaster scenario with a superimposed grid compared to a standard VR scene
24 (Chang et al. 2013), as Bles et al.'s (1998) subjective vertical conflict hypothesis would predict:
25 by providing a clear frame of reference for the visual vertical in VR, conflicts between
26 predicted and experienced subjective verticals are minimised.
27
28

29 Recent VR scenarios allow users to physically move in the VR environment, reducing
30 the conflict between visual and vestibular systems (Llorach et al. 2014). Despite the possibility
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

of cybersickness reduction through physical locomotion, limitations of physical space, particularly for home users of VR, may mean that users prefer to remain stationary and navigate the virtual environment by other means (for example, controllers) (Williams et al. 2007; Riecke et al. 2010). In addition, errors in position tracking and lag, while significantly improved in more recent HMDs, may also contribute to cybersickness development (Fiore et al. 2013; Kinsella et al. 2016; Palmisano et al. 2017).

Sensory conflicts can also be reduced by matching visual cues with artificial vestibular signals. Cevette et al. (2012) applied artificial vestibular stimulation (Galvanic Vestibular Stimulation) while participants used a flight simulator. Galvanic Vestibular Stimulation induced illusory **sensations of self-motion** which were purported to match the visual signals experienced by the participants. **Since the conflict between visual and vestibular signals was reduced**, a significant reduction in sickness symptoms was found. Similarly, Galvanic Vestibular Stimulation applied during turns in a driving simulator was suggested to reduce scores in motion sickness questionnaires and improve performance (Reed-Jones et al. 2007). In addition, Galvez-Garcia et al. (2015) found that applying Galvanic Cutaneous Stimulation either continuously or intermittently while participants used a driving simulator reduced sickness scores relative to a condition with no stimulation. These results suggest that the use of artificial stimulation may be a potential method for preventing cybersickness in VR.

VR after-effects: A re-adaptation to the real world

After-effects can arise following exposure to a variety of different sensory stimuli. One interesting and well-studied after-effect following exposure to passive motion, such as on a sea voyage, is so-called *mal de débarquement*. This syndrome induces illusory feelings of self-motion, such as bobbing or swaying lasting for days or even years (Van Ombergen et al. 2016).

1 Although less severe, after-effects can frequently develop in the hours and days following a
2 prolonged VR experience (Kellog et al. 1980; Gower and Fowlkes 1989). In one of the most
3
4 bizarre cases, a pilot had his view of the world invert 180 degrees while driving a car hours
5
6 after having been in a VR flight simulator (Kennedy et al. 1987). As a result of these after-
7
8 effects, many air force bases have mandatory policies which stipulate that pilots cannot fly an
9
10 aircraft up to 24 hours after exposure to a VR simulator. Also, many VR entertainment centres
11
12 require that users do not drive for several minutes after exposure. It is important to note that
13
14 some VR applications are designed specifically to carry over sensorimotor or behavioural
15
16 changes following exposure to the stimulus, for example for rehabilitation or training purposes
17
18 (Cameirao et al. 2011; Verschure 2011; Cameirao et al. 2012; Alaker et al. 2016; Pelargos et al.
19
20 2016; Valmaggia et al. 2016; Viñas-Diz and Sobrido-Prieto 2016). However, for the purposes
21
22 of this review we focus only on maladaptive after-effects.
23
24
25
26
27
28

29 After-effects might be induced by adaptation to conflicting sensory stimuli. One of the
30
31 most well-known examples is adaptation to prism lenses (Redding et al. 2005). On first
32
33 wearing the lenses, which displace the visual field, participants make frequent errors in pointing
34
35 and grasping. However, participants quickly adapt to the lenses and their accuracy increases.
36
37 On removal of the glasses, however, participants begin to make errors and a period of re-
38
39 adaptation is necessary before performance returns to normal (Clower et al. 1996). Adaptation
40
41 to VR may follow a similar pattern, however a thorough exploration of after-effects of VR
42
43 exposure has not yet been conducted.
44
45
46
47

48 Harm et al. (2008) found that proprioceptive coordination between the eyes, head and
49
50 hands was worse following 20 minutes of VR exposure. Critically, the performance was not
51
52 only immediately worse, but approached recovery only by 6 hours post-exposure. Similarly,
53
54 participants showed increased pointing errors immediately after 30 minutes of VR exposure
55
56 (Stanney et al. 1999).
57
58
59
60
61
62
63
64
65

Oculomotor after-effects have also been described after adaptation to VR. Di Girolamo and Pic (2001) found that vestibular-ocular reflex gain decreased immediately following 20 minutes of VR exposure, and took 30 minutes to return to baseline levels. Although preliminary, these results provide an insight into the potential complications of VR exposure.

A multisensory integration perspective for cybersickness

The evidence reviewed above suggests a pervasive influence of multisensory interaction in VR experience. In this section, we outline a conceptual scheme of how these could underlie cybersickness. Sensory inputs constantly reach the human brain. However, these signals need to be integrated to provide successful descriptions of the environment. Vestibular inputs make an essential contribution in this process, assessing whether visual signals are consistent or not with the movement and position of our head in space. Critically, when self-motion signals provided by the vestibular system cannot be aligned with those from visual cues, multisensory conflict occurs, potentially triggering cybersickness.

Under normal conditions, the visual and vestibular systems interact to provide information about self-motion. Optimal multisensory integration involves higher weighting of more reliable signals (Ernst and Banks, 2002). Thus, sensory cues are weighted according to their reliability, such that vestibular cues are given higher weighting when visual cues are unreliable and vice versa. A growing body of research has shown that visuo-vestibular integration is near-optimal for self-motion, with an overweighting of the vestibular cue relative to “true” optimality. Several results suggest that a process based on dynamic sensory reweighting may be important to explain cybersickness and adaptation to virtual environments (Butler et al. 2010; de Winkel et al. 2010; Fetsch et al. 2010; Angelaki et al. 2011). Three results are particularly relevant here. First, the contribution of optic flow to

perceived self-motion typically emerges only after several seconds of exposure to visual cues when inertial cues are not present, implying that a dynamic process of sensory re-weighting is necessary to resolve sensory conflicts before the perception of self-motion emerges (Young et al. 1973). Second, when vestibular and visual signals are conflicting, the optimal combination of cues cannot occur without a re-weighting of the original cues. In the real world, the vestibular cue is overweighted relative to optimality (Kaliuzhna et al. 2016, but see de Winkel et al. 2010). This implies that the sensory conflict experienced in virtual reality must be resolved by substantial down-weighting of vestibular cues and up-weighting of visual cues. Third, the results on visual or vestibular dominance in such conflicting conditions appear to vary across studies, suggesting that the human brain dynamically chooses which sensory cues are relevant for a particular situation (Young et al. 1973; Zacharias and Young 1981; Probst et al. 1985). However, taken together these results suggest that dynamic reweighting may be implicated in cybersickness.

Cybersickness may be due to the consequences of a conflict between the sensory signals which provide information about the body's orientation and motion. Visual signals tell users that they are moving in the environment. Since users are not actually moving, their vestibular architecture provides no corresponding cues of linear and angular acceleration. In view of this, when the perception of visual self-motion is not supported by inertial forces transmitted through the vestibular organs, a visuo-vestibular conflict is likely to occur, leading to cybersickness. Alteration of the weight of vestibular cues provides a means of resolving the visuo-vestibular conflict, reducing symptoms of cybersickness.

Multisensory theories suggest that multiple sensory signals need to be combined, and that the nervous system faces a key challenge in selecting the correct weighting for each signal in the combination. As in other cases of sensory conflict, resolution occurs by weighting each individual sensory signal according to its importance. In VR, the brain tends to habituate to

extract self-motion information from visual cues in a visuo-vestibular conflicting environment. Since vestibular information is usually highly reliable in determining the body's position and motion in space (Prsa et al. 2012; Kaliuzhna et al. 2016)², the vestibular signals must be down-weighted to avoid the occurrence of sensory conflicts. We suggest that a dynamic re-weighting function is an important element of cybersickness which has not yet been extensively researched. In VR, the dynamic re-weighting function increases the weight of visual signals about motion, and decreases the weight of vestibular information responsible for self-motion. As a result, the visuo-vestibular conflict is decreased and therefore no longer perceived, reducing cybersickness. Cybersickness symptoms typically develop within the first minutes of VR exposure (Stanney and Kennedy 1998; Davis et al. 2015). Thus, the re-weighting function must rapidly respond to the visuo-vestibular conflict to prevent the occurrence of symptoms. If the re-weighting function is slow to respond to the conflict, cybersickness may ensue. Interestingly, decreasing the reliability of vestibular cues by applying artificial noisy Galvanic Vestibular Stimulation has been shown to modulate vection perception (Weech and Troje 2017). This might be due to rapid re-weighting of visual and vestibular information. Critically, this approach can be implemented as a method to reduce cybersickness, as proposed by Weech and Troje (2017).

On return to the real world the individual is likely to move and explore the environment, causing a flow of both vestibular and visual signals about self-motion. After exposure to VR, the brain tends to habituate to extract self-motion information from visual cues in a vestibular-conflicting VR environment. As for traditional sensory adaptation phenomena, a form of negative correlation between a current percept and the adapted stimuli may take place (Barlow

² However, in some circumstances vestibular information can be misleading. For example, the semi-circular canals adapt to constant rotation and report no motion, while constant linear accelerations may be perceived as static tilts as in the somatogravic illusion – see Snyder, 1999, for further information.

and Hill 1963). An error correction process needs to occur in order to de-correlate the current percept from the adapted stimuli. Thus, a further re-weighting of both sensory signals is necessary for the optimal integration of sensory input. A period of re-adaptation must occur, whereby the vestibular cue must be up-weighted and visual cue down-weighted. During this period of re-adaptation, VR after-effects may occur over time until the vestibular cue is weighted to its usual state. The time-course of these after-effects is not yet fully known and may range from a period of minutes to hours or days (Kellog et al. 1980; Gower and Fowlkes 1989; Harm et al. 2008).

Traditionally, cybersickness has been explained by prior knowledge of predicted sensory consequences of self-motion (Reason and Brand 1975). Our conceptual scheme suggests a dynamic on-line re-weighting function of sensory cues for self-motion which can explain cybersickness, adaptation, and after-effects of VR. Reciprocal inhibitory vestibular-visual interactions support this hypothesis. Accordingly, PET studies using artificial vestibular stimulation demonstrated not only an activation of the PIVC but also a decrease in rCBF of the visual cortex (Wenzel et al. 1996; Brandt et al. 1998; Deutschländer et al. 2002). Similarly, Bense et al. (2001) showed deactivation of the occipital visual cortex induced by vestibular stimulation, and deactivation of the vestibular areas during optic flow.

Suggestions for future research

The conceptual scheme proposed here makes clear testable predictions about multisensory interactions in VR exposure and cybersickness. These could be investigated in lab-based cognitive experiments or in more applied VR scenarios. First, if the vestibular system plays a fundamental role in cybersickness, one might predict that synchronised passive movements or artificial vestibular stimulation may reduce the conflict between visual and

1 vestibular cues, preventing sickness and forgoing the need for sensory re-weighting function in
2 order to adapt to the VR environment. Indeed, recent reports (Reed-Jones et al. 2007; Cevette
3 et al. 2012; Galvez-Garcia et al. 2015) have demonstrated that both galvanic vestibular
4 stimulation and galvanic cutaneous stimulation can reduce sickness in simulators.
5
6
7

8
9 Second, if vestibular cues are down-weighted during VR exposure, one might expect
10 that physiological vestibular functioning is altered during or immediately after VR. Although
11 this aspect has not yet been investigated, one can imagine a testing battery investigating both
12 vestibular sensitivity and physiological functioning in order to identify whether the vestibular
13 organs adapt to VR.
14
15
16
17
18
19
20

21 Finally, individual variability in the re-weighting function may correlate with
22 cybersickness susceptibility. Those whose weighting function rapidly changes reliance on the
23 vestibular cue under VR conditions are potentially less likely to experience cybersickness, as
24 the cue conflict is rapidly resolved. By contrast, those with a higher reliance on the vestibular
25 cue under normal conditions may be more susceptible to cybersickness as the magnitude of the
26 cue conflict is greater. Extensive research on these predictions has yet to be conducted,
27 however we note that Balter et al. (2004) found no difference on sensory re-weighting abilities
28 between participants susceptible and non-susceptible to car-sickness. Participants susceptible
29 and non-susceptible to car-sickness were repeatedly administered with galvanic vestibular
30 stimulation while their body sway was measured. Although it was predicted that participants
31 who were less susceptible to car-sickness would habituate to the vestibular stimulation (and
32 therefore show reduced body sway) faster than participants who suffered from car-sickness,
33 both susceptible and non-susceptible participants showed similar habituation gains to the
34 vestibular stimulation. However, it is important to highlight that these findings do not refer to
35 conflicting stimuli; it is therefore possible that susceptible and non-susceptible individuals may
36 show differences in re-weighting when exposed to visual–vestibular conflicting signals.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Conclusion

While significant advances have been made in understanding cybersickness, there is broad scope for further advancement in this field. The symptom profile of cybersickness has been clearly delineated, as well as identification of factors which influence its development. However, explanations for why cybersickness occurs based on sensory conflict theory are tentative, and there is no identified predictive mechanism. In addition, a conspicuous gap in our knowledge concerns the after-effects of virtual reality exposure. A few studies have identified cybersickness symptoms, proprioceptive disruptions, postural instability and oculomotor symptoms as potential after-effects, however their time-course is unknown, despite the time-course of cybersickness itself being well-defined (Kennedy et al. 2000). The current sensory conflict models also cannot fully account for why these after-effects arise.

Here we argued that an approach based on multisensory integration could provide a predictive explanation for cybersickness, adaptation to VR and its after-effects. Under normal circumstances, visual and vestibular cues are optimally integrated for self-motion perception. However, in VR these cues conflict with one another, prompting cybersickness, and must be re-weighted with higher reliance on the visual cue to allow integration. This dynamic re-weighting function has the potential to explain adaptation and after-effects of VR, as well as individual differences in cybersickness susceptibility. It is hoped that this approach could further our knowledge of cybersickness, as well as lead to clearer avenues for prevention of cybersickness symptoms.

Acknowledgments

This work was supported by The British Academy award “Cybersickness: a perceptual information-processing approach” [grant number SG162313]. M.G. is further supported by a ESRC-DTC studentship.

References

- Akiduki, H., Nishiike, S., Watanabe, H., Matsuoka, K., Kubo, T. and Takeda, N. (2003) 'Visual-vestibular conflict induced by virtual reality in humans', *Neuroscience Letters*, 340(3), pp. 197–200. doi: 10.1016/S0304-3940(03)00098-3.
- Alaker, M., Wynn, G. R. and Arulampalam, T. (2016) 'Virtual reality training in laparoscopic surgery: A systematic review & meta-analysis', *International Journal of Surgery*, 29, pp. 85–94. doi: 10.1016/j.ijssu.2016.03.034.
- Angelaki, D. E., Gu, Y. and Deangelis, G. C. (2011) 'Visual and vestibular cue integration for heading perception in extrastriate visual cortex.', *The Journal of physiology*, 589(Pt 4), pp. 825–833. doi: 10.1113/jphysiol.2010.194720.
- Arns, L. L. and Cerney, M. M. (2005) 'The relationship between age and incidence of cybersickness among immersive environment users', *IEEE Proceedings. VR 2005. Virtual Reality, 2005.*, 2005, pp. 267–268. doi: 10.1109/VR.2005.1492788.
- Ash, A. and Palmisano, S. (2012) 'Vection during conflicting multisensory information about the axis, magnitude, and direction of self-motion', *Perception*, 41(3), pp. 253–267. doi: 10.1068/p7129.
- Ash, A., Palmisano, S. and Kim, J. (2011) 'Vection in depth during consistent and inconsistent multisensory stimulation', *Perception*, 40(2), pp. 155–174. doi: 10.1068/p6837.
- Azmandian, M., Hancock, M., Benko, H., Ofek, E. and Wilson, A. D. (2016) 'Haptic Retargeting: Dynamic Repurposing of Passive Haptics for Enhanced Virtual Reality Experiences', *Proceedings of the 2016 CHI Conference on Human Factors in Computing Systems - CHI '16*, pp. 1968–1979. doi: 10.1145/2858036.2858226.
- Balter, S. G. T., Stokroos, R. J., Van De Laar, M. M. M., Hendrice, N. and Kingma, H. (2004) 'Habituation to Galvanic Vestibular Stimulation for Analysis of Susceptibility to

- Carsickness', *Acta Oto-Laryngologica*, 124(6), pp. 690–694. doi: 10.1080/00016480410017242.
- Barlow, H. B. and Hill, R. M. (1963) 'Evidence for a physiological explanation of the waterfall phenomenon and figural after-effects.', *Nature*, 200, pp. 1345–7.
- Barra, J., Marquer, A., Joassin, R., Reymond, C., Metge, L., Chauvineau, V. and Pérennou, D. (2010) 'Humans use internal models to construct and update a sense of verticality', *Brain*, 133(12), pp. 3552–3563. doi: 10.1093/brain/awq311.
- Barrett, J. (2004) 'Side effects of virtual environments: A review of the literature', *Information Sciences*. Available at: <http://oai.dtic.mil/oai/oai?verb=getRecord&metadataPrefix=html&identifier=ADA426109>.
- Bastug, E. et al., 2017. Toward Interconnected Virtual Reality: Opportunities, Challenges, and Enablers. *IEEE Communications Magazine*, 55(6), pp.110–117.
- Bellini, H., Chen, W., Sugiyama, M., Shin, M., Alam, S. and Takayama, D. (2016) 'Virtual & Augmented Reality: Understanding the race for the next computing platform', *Profiles in Innovation*. doi: 10.1017/CBO9781107415324.004.
- Bense, S., Stephan, T., Yousry, T. A., Brandt, T. and Dieterich, M. (2001) 'Multisensory cortical signal increases and decreases during vestibular galvanic stimulation (fMRI).', *Journal of neurophysiology*, 85(2), pp. 886–99.
- Berthoz, A., Israël, I., Georges-François, P., Grasso, R. and Tsuzuku, T. (1995) 'Spatial memory of body linear displacement: what is being stored?', *Science (New York, N.Y.)*, 269(5220), pp. 95–8.
- Bles, W., Bos, J. E., De Graaf, B., Groen, E. and Wertheim, A. H. (1998) 'Motion sickness: Only one provocative conflict?', *Brain Research Bulletin*, 47(5), pp. 481–487. doi: 10.1016/S0361-9230(98)00115-4.

- Bonato, F., Bubka, A. and Palmisano, S. (2009) 'Combined pitch and roll and cybersickness in a virtual environment', *Aviation Space and Environmental Medicine*, 80(11), pp. 941–945. doi: 10.3357/ASEM.2394.2009.
- Bos, J. E., Bles, W. and Groen, E. L. (2008) 'A theory on visually induced motion sickness', *Displays*, 29(2), pp. 47–57. doi: 10.1016/j.displa.2007.09.002.
- Brandt, T., Bartenstein, P., Janek, A. and Dieterich, M. (1998) 'Reciprocal inhibitory visual-vestibular interaction. Visual motion stimulation deactivates the parieto-insular vestibular cortex', *Brain*, 121(9), pp. 1749–1758. doi: 10.1093/brain/121.9.1749.
- Bremmer, F., Klam, F., Duhamel, J.-R., Ben Hamed, S. and Graf, W. (2002) 'Visual-vestibular interactive responses in the macaque ventral intraparietal area (VIP).', *The European journal of neuroscience*, 16(8), pp. 1569–86.
- Butler, J. S., Smith, S. T., Campos, J. L. and Bühlhoff, H. H. (2010) 'Bayesian integration of visual and vestibular signals for heading.', *Journal of vision*, 10(11), p. 23. doi: 10.1167/10.11.23.
- Cameirão, M. S., Badia, S. B. I., Duarte, E., Frisoli, A. and Verschure, P. F. M. J. (2012) 'The combined impact of virtual reality neurorehabilitation and its interfaces on upper extremity functional recovery in patients with chronic stroke', *Stroke*, 43(10), pp. 2720–2728. doi: 10.1161/STROKEAHA.112.653196.
- Cevette, M. J., Stepanek, J., Cocco, D., Galea, A. M., Pradhan, G. N., Wagner, L. S., Oakley, S. R., Smith, B. E., Zapala, D. A. and Brookler, K. H. (2012) 'Oculo-Vestibular Recoupling Using Galvanic Vestibular Stimulation to Mitigate Simulator Sickness', *Aviation Space and Environmental Medicine*, 83(6), pp. 549–555. doi: 10.3357/ASEM.3239.2012.
- Chance, S. S., Gaunet, F., Beall, A. C. and Loomis, J. M. (1998) 'Locomotion Mode Affects the Updating of Objects Encountered During Travel: The Contribution of Vestibular and

- Proprioceptive Inputs to Path Integration', *Presence: Teleoperators and Virtual Environments*, 7(2), pp. 168–178. doi: 10.1162/105474698565659.
- Chang, E., Hwang, I., Jeon, H., Chun, Y., Kim, H. T. and Park, C. (2013) 'Effects of rest frames on cybersickness and oscillatory brain activity', *2013 International Winter Workshop on Brain-Computer Interface, BCI 2013*, pp. 62–64. doi: 10.1109/IWW-BCI.2013.6506631.
- Chen, A., DeAngelis, G. C. and Angelaki, D. E. (2010) 'Macaque parieto-insular vestibular cortex: responses to self-motion and optic flow.', *The Journal of neuroscience : the official journal of the Society for Neuroscience*. NIH Public Access, 30(8), pp. 3022–42. doi: 10.1523/JNEUROSCI.4029-09.2010.
- Chen, D. J., Bao, B., Zhao, Y. and So, R. H. Y. (2015) 'Visually induced motion sickness when viewing visual oscillations of different frequencies along the fore-and-aft axis: keeping velocity versus amplitude constant.', *Ergonomics*, 139(July), pp. 1–9. doi: 10.1080/00140139.2015.1078501.
- Chen, W., Chao, J. G., Chen, X. W., Wang, J. K. and Tan, C. (2015) 'Quantitative orientation preference and susceptibility to space motion sickness simulated in a virtual reality environment', *Brain Research Bulletin*. Elsevier Inc., 113, pp. 17–26. doi: 10.1016/j.brainresbull.2015.01.007.
- Cheung, B. S., Howard, I. P. and Money, K. E. (1991) 'Visually-induced sickness in normal and bilaterally labyrinthine-defective subjects', *Aviation Space and Environmental Medicine*, 62(6), pp. 527–531.
- Clemes, S. A. and Howarth, P. A. (2005) 'The Menstrual Cycle and Susceptibility to Virtual Simulation Sickness', *Journal of Biological Rhythms*, 20(1), pp. 71–82. doi: 10.1177/0748730404272567.

- Clower, D. M., Hoffman, J. M., Votaw, J. R., Faber, T. L., Woods, R. P. and Alexander, G. E. (1996) 'Role of posterior parietal cortex in the recalibration of visually guided reaching', *Nature*, pp. 618–621. doi: 10.1038/383618a0.
- Cobb, S. V. G., Nichols, S., Ramsey, A. and Wilson, J. R. (1999) 'Virtual Reality-Induced Symptoms and Effects (VRISE)', *Presence*, 8(2), pp. 169–186. doi: 10.1162/105474699566152.
- Davis, S., Nesbitt, K. and Nalivaiko, E. (2014) 'A Systematic Review of Cybersickness', *Proceedings of the 2014 Conference on Interactive Entertainment - IE2014*, pp. 1–9. doi: 10.1145/2677758.2677780.
- Davis, S., Nesbitt, K. and Nalivaiko, E. (2015) 'Comparing the onset of cybersickness using the Oculus Rift and two virtual roller coasters', *11th Australasian Conference on Interactive Entertainment (IE 2015)*, (January), pp. 27–30. doi: 10.17973/MMSJ.2015.
- Dennison, M. S. and D'Zmura, M. (2017) 'Cybersickness without the wobble: Experimental results speak against postural instability theory', *Applied Ergonomics*. Elsevier Ltd, 58, pp. 215–223. doi: 10.1016/j.apergo.2016.06.014.
- Dennison, M. S., Wisti, A. Z. and D'Zmura, M. (2016) 'Use of physiological signals to predict cybersickness', *Displays*. Elsevier B.V., 44, pp. 42–52. doi: 10.1016/j.displa.2016.07.002.
- Deutschländer, A., Bense, S., Stephan, T., Schwaiger, M., Brandt, T. and Dieterich, M. (2002) 'Sensory system interactions during simultaneous vestibular and visual stimulation in PET.', *Human brain mapping*, 16(2), pp. 92–103.
- Di Girolamo, S. and Pic, P. (2001) 'Vestibulo-Ocular Reflex Modification after Virtual Environment Exposure', *Acta Oto-Laryngologica*, 121(2), pp. 211–215. doi: 10.1080/000164801300043541.

- Ernst, M. O. and Banks, M. S. (2002) ‘Humans integrate visual and haptic information in a statistically optimal fashion.’, *Nature*, 415(6870), pp. 429–433. doi: 10.1038/415429a.
- Ernst, M. O. and Bühlhoff, H. H. (2004) ‘Merging the senses into a robust percept’, *Trends in Cognitive Sciences*, 8(4), pp. 162–169. doi: 10.1016/j.tics.2004.02.002.
- Fasold, O., von Brevern, M., Kuhberg, M., Ploner, C. J., Villringer, A., Lempert, T. and Wenzel, R. (2002) ‘Human vestibular cortex as identified with caloric stimulation in functional magnetic resonance imaging.’, *NeuroImage*, 17(3), pp. 1384–1393. doi: S1053811902912413 [pii].
- Ferrè, E. R. and Haggard, P. (2015) ‘Vestibular–Somatosensory Interactions: A Mechanism in Search of a Function?’, *Multisensory Research*, 28(5–6), pp. 559–579.
- Fetsch, C. R., Deangelis, G. C. and Angelaki, D. E. (2010) ‘Visual-vestibular cue integration for heading perception: Applications of optimal cue integration theory’, *European Journal of Neuroscience*, 31(10), pp. 1721–1729. doi: 10.1038/jid.2014.371.
- Fetsch, C. R., Turner, A. H., DeAngelis, G. C. and Angelaki, D. E. (2009) ‘Dynamic reweighting of visual and vestibular cues during self-motion perception.’, *The Journal of neuroscience: the official journal of the Society for Neuroscience*, 29(49), pp. 15601–15612. doi: 10.1523/JNEUROSCI.2574-09.2009.
- Fiore, L. P., Coben, E., Merritt, S., Liu, P. and Interrante, V. (2013) ‘Towards Enabling More Effective Locomotion in VR Using a Wheelchair-based Motion Platform’, *Proceedings of the 5th Joint Virtual Reality Conference*, pp. 83–90. doi: 10.2312/EGVE.JVRC13.083-090.
- Flanagan, M. B., May, J. G. and Dobie, T. G. (2005) ‘Sex Differences in Tolerance to Visually-Induced Motion Sickness’, *Aviation, Space, and Environmental Medicine*, 76(7), pp. 642–646.

- Galvez-Garcia, G., Hay, M. and Gabaude, C. (2015) 'Alleviating Simulator Sickness with Galvanic Cutaneous Stimulation', *Human Factors: The Journal of the Human Factors and Ergonomics Society*, 57(4), pp. 649–657. doi: 10.1177/0018720814554948.
- Gibson, J. J. (1950) 'The Perception of Visual Surfaces', *The American Journal of Psychology*, 63(3), pp. 367–384.
- Golding, J. (2016) *Motion sickness*. 1st edn, *Handbook of clinical neurology*. 1st edn. Elsevier B.V. doi: 10.1016/B978-0-444-63437-5.00027-3.
- Golding, J. F., Kadzere, P. and Gresty, M. A. (2005) 'Motion Sickness Susceptibility Fluctuates Through the Menstrual Cycle', *Aviation, Space, and Environmental Medicine*, 76(10), pp. 970–973. doi: 10.1196/annals.1429.018.
- Gower, D. W. and Fowlkes, J. E. (1989) 'Simulator Sickness in the UH-60 (Black Hawk) Flight Simulator.', *USAARL 89-20 (AD-A214 434), U.S. Army Aeromedical Research Laboratory*.
- Green, A. M. and Angelaki, D. E. (2010) 'Multisensory integration: Resolving sensory ambiguities to build novel representations', *Current Opinion in Neurobiology*. Elsevier Ltd, 20(3), pp. 353–360. doi: 10.1016/j.conb.2010.04.009.
- Greenlee, M. W., Frank, S. M., Kaliuzhna, M., Blanke, O., Bremmer, F., Churan, J., Cuturi, L. F., MacNeilage, P. R. and Smith, A. T. (2016) 'Multisensory Integration in Self Motion Perception', *Multisensory Research*, 29(March 2016), pp. 1–32. doi: 10.1163/22134808-00002527.
- Greybeil, A. (1970) 'Susceptibility to acute motion sickness in blind persons.', *Aerospace Medicine*, 41(6), pp. 650–653.
- Gu, Y., Angelaki, D. E. and DeAngelis, G. C. (2008) 'Neural correlates of multisensory cue integration in macaque MSTd', *Nature Neuroscience*, 11(10), pp. 1201–1210. doi: 10.1038/nn.2191.

- Guldin, W. O. and Grüsser, O. J. (1998) 'Is there a vestibular cortex?', *Trends in Neurosciences*, 21(6), pp. 254–259. doi: 10.1016/S0166-2236(97)01211-3.
- Han, K., Park, C., Kim, E., Kim, D., Woo, S., Jeong, J., Hwang, I. and Kim, H. (2011) 'Effects of Different Types of 3D Rest Frames on Reducing Cybersickness in a Virtual Environment', *i-Perception*. SAGE PublicationsSage UK: London, England, 2(8), pp. 861–861. doi: 10.1068/ic861.
- Harm, D. L., Taylor, L. C., Reschke, M. F., Somers, J. T. and Bloomberg, J. J. (2008) 'Sensorimotor coordination aftereffects of exposure to a virtual environment', *Visual Computer*, 24(11), pp. 995–999. doi: 10.1007/s00371-008-0277-1.
- Harsora, J., Khanvilkar, A., Sayyad, M. and Road, M. (2017) 'A Systematic Literature Review On Virtual Reality- The Oculus Rift', *International Journal of Research in Science and Engineering*, (7), pp. 1–9.
- Herbelin, B., Salomon, R., Serino, A. and Blanke, O. (2015) '5. Neural Mechanisms of Bodily Self-Consciousness and the Experience of Presence in Virtual Reality', *Human Computer Confluence*, pp. 80–96. doi: 10.1515/9783110471137-005.
- Hill, K. J. and Howarth, P. A. (2000) 'Habituation to the side effects of immersion in a virtual environment', *Displays*, 21(1), pp. 25–30. doi: 10.1016/S0141-9382(00)00029-9.
- Howarth, P. A. and Hodder, S. G. (2008) 'Characteristics of habituation to motion in a virtual environment', *Displays*, 29(2), pp. 117–123. doi: 10.1016/j.displa.2007.09.009.
- Hsu, C. et al., 2017. Is Foveated Rendering Perceivable in Virtual Reality? Exploring the Efficiency and Consistency of ality Assessment Methods. , pp.55–63. Available at: http://dirl.iis.sinica.edu.tw/pub/hsu17_is_foveated_rendering_perceivable.pdf.
- Israël, I. and Berthoz, A. (1989) 'Contribution of the otoliths to the calculation of linear displacement.', *Journal of neurophysiology*, 62(1), pp. 247–63. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2754476> (Accessed: 5 October 2017).

- Israël, I., Chapuis, N., Glasauer, S., Charade, O. and Berthoz, A. (1993) 'Estimation of passive horizontal linear whole-body displacement in humans.', *Journal of neurophysiology*, 70(3), pp. 1270–3. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8229174> (Accessed: 5 October 2017).
- Jürgens, R., Kliegl, K., Kassubek, J., & Becker, W. (2016). Optokinetic circularvection: a test of visual–vestibular conflict models ofvection nascensy. *Experimental Brain Research*, 234(1), 67–81. <https://doi.org/10.1007/s00221-015-4433-3>
- Kaliuzhna, M., Ferrè, E. R., Herbelin, B., Blanke, O. and Haggard, P. (2016) 'Multisensory effects on somatosensation: a trimodal visuo-vestibular-tactile interaction', *Scientific Reports*, 6.
- Kato, K. and Kitazaki, S. (2008) 'Improvement of Ease of Viewing Images on an In-vehicle Display and Reduction of Carsickness', *Human Factors in Driving, Seating Comfort and Automotive Telematics*. doi: 10.4271/2008-01-0565.
- Kellog, R. S., Castore, C. and Coward, R. (1980) 'Psychophysiological Effects of Training in a Full Vision Simulator.', *Annual Scientific Meeting of the Aerospace Medical Association*.
- Kennedy, R. S., Berbaum, K. S., Lilienthal, M. G., Dunlap, W. P., Mulligan, B. E. and Funaro, J. F. (1987) 'Guidelines for Alleviation of Simulator Sickness Symptomatology', pp. 559–563. doi: 10.1111/j.1559-3584.1933.tb04295.x.
- Kennedy, R. S., Drexler, J. and Kennedy, R. C. (2010) 'Research in visually induced motion sickness', *Applied Ergonomics*, 41(4), pp. 494–503. doi: 10.1016/j.apergo.2009.11.006.
- Kennedy, R. S., Drexler, J. M., Compton, D. E., Stanney, K. M., Lanham, D. S. and Harm, D. L. (2003) 'Configural scoring of simulator sickness, cybersickness and space adaptation syndrome: similarities and differences', *Virtual and Adaptive Environments*

Applications Implications and Human Performance Issues, pp. 247–278. doi:
10.1201/9781410608888.ch12.

Kennedy, R. S., Lane, N. E., Berbaum, K. S. and Lilienthal, M. G. (1993) ‘Simulator Sickness Questionnaire: An Enhanced Method for Quantifying Simulator Sickness’, *The International Journal of Aviation Psychology*, 3(3), pp. 203–220.

Kennedy, R. S., Stanney, K. M., & Dunlap, W. P. (2000). ‘Duration and exposure to virtual environments: sickness curves during and across sessions’. *Presence: Teleoperators and virtual environments*, 9(5), 463-472.

Keshavarz, B. (2013) ‘Exploring Behavioral Methods to Reduce Visually Induced Motion Sickness in Virtual Environments’, *5th International Conference, VAMR 2013, Held as Part of HCI International 2013*, 8021, p. 399. doi: 10.1007/978-3-642-39405-8.

Keshavarz, B. and Hecht, H. (2011) ‘Axis rotation and visually induced motion sickness: The role of combined roll, pitch, and yaw motion’, *Aviation Space and Environmental Medicine*, 82(11), pp. 1023–1029. doi: 10.3357/ASEM.3078.2011.

Keshavarz, B., Riecke, B. E., Hettinger, L. J. and Campos, J. L. (2015) ‘Vection and visually induced motion sickness: How are they related?’, *Frontiers in Psychology*, 6(APR), pp. 1–11. doi: 10.3389/fpsyg.2015.00472.

Kim, Y. Y., Kim, E. N., Park, M. J., Park, K. S., Ko, H. D. and Kim, H. T. (2008) ‘The Application of Biosignal Feedback for Reducing Cybersickness from Exposure to a Virtual Environment’, *Presence: Teleoperators and Virtual Environments*, 17(1), pp. 1–16. doi: 10.1162/pres.17.1.1.

Kim, Y. Y., Kim, H. T. J., Kim, E. N., Ko, H. D. and Kim, H. T. J. (2005) ‘Characteristic changes in the physiological components of cybersickness’, *Psychophysiology*, 42(5), pp. 616–625. doi: 10.1007/s00234-005-1388-2.

- Kim, Y. Y., Kim, H. T. J., Ko, H. D. and Kim, H. T. J. (2001) 'Psychophysiological changes by navigation in a virtual reality', *Annual Reports of the Research Reactor Institute, Kyoto University*, 4, pp. 3773–3776. doi: 10.1109/IEMBS.2001.1019659.
- Kinsella, A., Mattfeld, R., Muth, E. and Hoover, A. (2016) 'Frequency, Not Amplitude, of Latency Affects Subjective Sickness in a Head-Mounted Display', *Aerospace Medicine and Human Performance*, 87(7), pp. 604–609. doi: 10.3357/AMHP.4351.2016.
- Knill, D. C. and Pouget, A. (2004) 'The Bayesian brain: The role of uncertainty in neural coding and computation', *Trends in Neurosciences*, 27(12), pp. 712–719. doi: 10.1016/j.tins.2004.10.007.
- Koslucher, F., Haaland, E. and Stoffregen, T. A. (2016) 'Sex differences in visual performance and postural sway precede sex differences in visually induced motion sickness', *Experimental Brain Research*. Springer Berlin Heidelberg, 234(1), pp. 313–322. doi: 10.1007/s00221-015-4462-y.
- Kushner, D. (2016) 'Will Virtual Reality Change Your Life?', *Rolling Stone*, May. Available at: <http://www.rollingstone.com/culture/features/will-virtual-reality-change-your-life-20160523> (Accessed: 10 July 2017).
- Lackner, J. R. (2014) 'Motion sickness: More than nausea and vomiting', *Experimental Brain Research*, 232(8), pp. 2493–2510. doi: 10.1007/s00221-014-4008-8.
- LaViola, J. J. (2000) 'A discussion of cybersickness in virtual environments', *ACM SIGCHI Bulletin*, 32(1), pp. 47–56. doi: 10.1145/333329.333344.
- Lee, W.-T. et al., 2017. High-resolution 360 Video Foveated Stitching for Real-time VR. *Computer Graphics Forum*, 36(7), pp.115–123. Available at: <http://doi.wiley.com/10.1111/cgf.13277>.

Liu, C. L. (2014) 'A study of detecting and combating cybersickness with fuzzy control for the elderly within 3D virtual stores', *International Journal of Human Computer Studies*. Elsevier, 72(12), pp. 796–804. doi: 10.1016/j.ijhcs.2014.07.002.

Llorach, G., Evans, A. and Blat, J. (2014) 'Simulator sickness and presence using HMDs: comparing use of a game controller and a position estimation system', *20th ACM Symposium on Virtual Reality Software and Technology - VRST '14*, pp. 137–140. doi: 10.1145/2671015.2671120.

Lo, W. T. and So, R. H. Y. (2001) 'Cybersickness in the presence of scene rotational movements along different axes', *Applied Ergonomics*, pp. 1–14. doi: 10.1016/S0003-6870(00)00059-4.

Lopez, C. and Blanke, O. (2011) 'The thalamocortical vestibular system in animals and humans', *Brain Research Reviews*. Elsevier B.V., 67(1–2), pp. 119–146. doi: 10.1016/j.brainresrev.2010.12.002.

Lopez, C., Blanke, O. and Mast, F. W. (2012) 'The human vestibular cortex revealed by coordinate-based activation likelihood estimation meta-analysis', *Neuroscience*, 212, pp. 159–179. doi: 10.1016/j.neuroscience.2012.03.028.

Merhi, O., Faugloire, E., Flanagan, M. and Stoffregen, T. A. (2007) 'Motion Sickness, Console Video Games, and Head-Mounted Displays', *Human Factors*, 49(5), pp. 920–934. doi: 10.1518/001872007X230262.

Moss, J. D., Austin, J., Salley, J., Coats, J., Williams, K. and Muth, E. R. (2011) 'The effects of display delay on simulator sickness', *Displays*, 32(4), pp. 159–168. doi: 10.1016/j.displa.2011.05.010.

Munafo, J., Diedrick, M. and Stoffregen, T. A. (2017) 'The virtual reality head-mounted display Oculus Rift induces motion sickness and is sexist in its effects', *Experimental*

Brain Research. Springer Berlin Heidelberg, 235(3), pp. 889–901. doi: 10.1007/s00221-016-4846-7.

Nalivaiko, E., Davis, S. L., Blackmore, K. L., Vakulin, A. and Nesbitt, K. V. (2015) ‘Cybersickness provoked by head-mounted display affects cutaneous vascular tone, heart rate and reaction time’, *Physiology and Behavior*. Elsevier Inc., 151, pp. 583–590. doi: 10.1016/j.physbeh.2015.08.043.

Nichols, S. (2000) ‘Individual characteristics and experiences of virtual reality induced symptoms and effects’, *XIV Triennial Congress of the International Ergonomics Association and 44th Annual Meeting of the Human Factors and Ergonomics Society*, 1, pp. 538–541. doi: 10.1177/154193120004400514.

Nishiike, S., Okazaki, S., Watanabe, H., Akizuki, H., Imai, T., Uno, A., Kitahara, T., Horii, A., Takeda, N. and Inohara, H. (2013) ‘The effect of visual-vestibulosomatosensory conflict induced by virtual reality on postural stability in humans’, *The journal of medical investigation : JMI*, 60(3–4), pp. 236–239. doi: 10.2152/jmi.60.236.

Oman, C. M. (1988) ‘Motion Sickness: a synthesis and evaluation of the sensory conflict theory’, *Canadian Journal of Physiology Pharmacology*, 68, pp. 294–303.

Oman, C. M. (2012) ‘Are evolutionary hypotheses for motion sickness “just-so” stories?’, *Journal of Vestibular Research: Equilibrium and Orientation*, 22(2–3), pp. 117–127. doi: 10.3233/VES-2011-0432.

Padmanaban, N. et al., 2017. Optimizing virtual reality for all users through gaze-contingent and adaptive focus displays. *Proceedings of the National Academy of Sciences of the United States of America*, 114(9), pp.2183–2188. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/28193871>

Paillard, A. C., Quarck, G., Paolino, F., Denise, P., Paolino, M., Golding, J. F. and Ghulyan-Bedikian, V. (2013) ‘Motion sickness susceptibility in healthy subjects and vestibular

- patients: Effects of gender, age and trait-anxiety', *Journal of Vestibular Research: Equilibrium and Orientation*, 23(4–5), pp. 203–210. doi: 10.3233/VES-130501.
- Palmisano, S., Allison, R. S., Schira, M. M. and Barry, R. J. (2015) 'Future challenges for vection research: Definitions, functional significance, measures, and neural bases', *Frontiers in Psychology*, 6(FEB), pp. 1–15. doi: 10.3389/fpsyg.2015.00193.
- Palmisano, S., Mursic, R. and Kim, J. (2017) 'Vection and cybersickness generated by head-and-display motion in the Oculus Rift', *Displays*, 46, pp. 1–8. doi: 10.1016/j.displa.2016.11.001.
- Peck, T. C., Fuchs, H. and Whitton, M. C. (2011) 'An evaluation of navigational ability comparing Redirected Free Exploration with Distractors to Walking-in-Place and joystick locomotion interfaces', *Proceedings - IEEE Virtual Reality*, pp. 55–62. doi: 10.1109/VR.2011.5759437.
- Pelargos, P. E., Nagasawa, D. T., Lagman, C., Tenn, S., Demos, J. V., Lee, S. J., Bui, T. T., Barnette, N. E., Bhatt, N. S., Ung, N., Bari, A., Martin, N. A. and Yang, I. (2016) 'Utilizing virtual and augmented reality for educational and clinical enhancements in neurosurgery', *Journal of Clinical Neuroscience*. Elsevier Ltd, 35((in press)), pp. 1–4. doi: 10.1016/j.jocn.2016.09.002.
- Probst, T., Straube, A. and Bles, W. (1985) 'Differential Effects Of Ambivalent Visual-Vestibular-Somatosensory Stimulation On The Perception Of Self-Motion', *Behavioural Brain Research*, 16(1), pp. 71–79. doi: 10.1016/0166-4328(85)90083-X.
- Prsa, M., Gale, S. and Blanke, O. (2012) 'Self-motion leads to mandatory cue fusion across sensory modalities', *Journal of Neurophysiology*, 108(8), pp. 2282–2291. doi: 10.1152/jn.00439.2012.
- Reason, J. T. (1978) 'Motion sickness adaptation: a neural mismatch model.', *Journal of the Royal Society of Medicine*, 71(11), pp. 819–29. doi: 10.1177/014107687807101109.

- Reason, J. T. and Brand, J. J. (1975) *Motion Sickness*. New York, NY, NY: Academic Press.
- Rebenitsch, L. and Owen, C. (2014) 'Individual variation in susceptibility to cybersickness', *Proceedings of the 27th annual ACM symposium on User interface software and technology*, pp. 309–317. doi: 10.1145/2642918.2647394.
- Rebenitsch, L. and Owen, C. (2016) 'Review on cybersickness in applications and visual displays', *Virtual Reality*. Springer London, 20(2), pp. 101–125. doi: 10.1007/s10055-016-0285-9.
- Redding, G. M., Rossetti, Y. and Wallace, B. (2005) 'Applications of prism adaptation: a tutorial in theory and method.', *Neuroscience Biobehavioural Review*, 29(3), pp. 431–444.
- Reed-Jones, R. J., Reed-Jones, J. G., Trick, L. M. and Vallis, L. a (2007) 'Can galvanic vestibular stimulation reduce simulator adaptation syndrome?', *Human Factors*, (June 2014), pp. 534–540. Available at: http://www.psychology.uoguelph.ca/faculty/trick/Documents/Articles/086_ReedJonesTrick.pdf.
- Regan, E. C. (1995) 'Some evidence of adaptation to immersion in virtual reality', *Displays*, 16(3), pp. 135–139. doi: 10.1016/0141-9382(96)81213-3.
- Riccio, G. E. and Stoffregen, T. A. (1991) 'An ecological theory of motion sickness and postural instability', *Ecological Psychology*, 3(3), pp. 195–240.
- Riecke, B. E., Bodenheimer, B., McNamara, T. P., Williams, B., Peng, P. and Feuereissen, D. (2010) 'Do we need to walk for effective virtual reality navigation? Physical rotations alone may suffice', *Lecture Notes in Computer Science (including subseries Lecture Notes in Artificial Intelligence and Lecture Notes in Bioinformatics)*, 6222 LNAI, pp. 234–247. doi: 10.1007/978-3-642-14749-4_21.

1 Sharples, S., Cobb, S., Moody, A. and Wilson, J. R. (2008) 'Virtual reality induced symptoms
2 and effects (VRISE): Comparison of head mounted display (HMD), desktop and
3 projection display systems', *Displays*, 29(2), pp. 58–69. doi:
4 10.1016/j.displa.2007.09.005.
5
6
7
8

9 da Silva Cameirão, M., Bermúdez Badia, S., Duarte, E. and FMJ Verschure, P. (2011) 'Virtual
10 reality based rehabilitation speeds up functional recovery of the upper extremities after
11 stroke: A randomized controlled pilot study in the acute phase of stroke using the
12 Rehabilitation Gaming System', *Restorative Neurology and Neuroscience*. IOS Press,
13 29, pp. 287–298. doi: 10.3233/RNN-2011-0599.
14
15
16
17
18
19
20
21

22 Smart, L. J., Stoffregen, T. A. and Benoit, B. G. (2002) 'Visually Induced Motion Sickness
23 Predicted by Postural Instability', *Human Factors: The Journal of the Human Factors
24 and Ergonomics Society*, 44(3), pp. 451–465. doi: 10.1518/0018720024497745.
25
26
27
28

29 Snyder, L. (1999) 'This way up: illusions and internal models in the vestibular system', *Nature
30 Neuroscience*, 2(5), pp. 396–398. doi: 10.1038/8056.
31
32
33

34 So, R. H. Y. and Lo, W. T. (1999) 'Cybersickness: an experimental study to isolate the effects
35 of rotational scene oscillations', *Proceedings IEEE Virtual Reality (Cat. No.
36 99CB36316)*, pp. 237–241. doi: 10.1109/VR.1999.756957.
37
38
39
40

41 Stanney, K. M., Hale, K. S., Nahmens, I. and Kennedy, R. S. (2003) 'What to expect from
42 immersive virtual environment exposure: influences of gender, body mass index, and
43 past experience.', *Human factors*, 45(3), pp. 504–520. doi:
44 10.1518/hfes.45.3.504.27254.
45
46
47
48
49
50

51 Stanney, K. M. and Kennedy, R. S. (1998) 'Aftereffects from Virtual Environment Exposure:
52 How Long do They Last?', *Proceedings of the Human Factors and Ergonomics Society
53 Annual Meeting*, 42(21), pp. 1476–1480. doi: 10.1177/154193129804202103.
54
55
56
57
58
59
60
61
62
63
64
65

- Stanney, K. M., Kennedy, R. S. and Drexler, J. M. (1997) 'Cybersickness is not simulator sickness', *Proceedings of the Human Factors and Ergonomics Society 41st Annual Meeting*, pp. 1138–1142. doi: 10.1177/107118139704100292.
- Stanney, K. M., Kennedy, R. S., Drexler, J. M. and Harm, D. L. (1999) 'Motion sickness and proprioceptive aftereffects following virtual environment exposure', *Applied Ergonomics*, 30(1), pp. 27–38. doi: 10.1016/S0003-6870(98)00039-8.
- Stanney, K. M., Kingdon, K. S., Graeber, D. and Kennedy, R. S. (2002) 'Human Performance in Immersive Virtual Environments: Human Performance in Immersive Virtual Environments: Effects of Exposure Duration, User Control, and Scene Complexity', *Human Performance*, 15(4), pp. 339–366. doi: 10.1207/S15327043HUP1504.
- Stein, B. E., London, N., Wilkinson, L. K. and Price, D. D. (1996) 'Enhancement of Perceived Visual Intensity by Auditory Stimuli: A Psychophysical Analysis', *Journal of Cognitive Neuroscience*, 8(6), pp. 497–506.
- Stoffregen, T. A., Hettinger, L. J., Haas, M. W., Roe, M. M. and Smart, L. J. (2000) 'Postural Instability and Motion Sickness in a Fixed-Base Flight Simulator', *Human Factors: The Journal of the Human Factors and Ergonomics Society*, 42(3), pp. 458–469. doi: 10.1518/001872000779698097.
- Stoffregen, T. A. and Smart, L. J. (1998) 'Postural instability precedes motion sickness', *Brain Research Bulletin*, 47(5), pp. 437–448. doi: 10.1016/S0361-9230(98)00102-6.
- Sutherland, I. E. (1968) 'A Head-mounted Three Dimensional Display', in *Proceedings of the December 9-11, 1968, Fall Joint Computer Conference, Part I. New York, NY, USA: ACM (AFIPS '68 (Fall, part I))*, pp. 757–764. doi: 10.1145/1476589.1476686**
- Treisman, M. (1977) 'Motion Sickness: An Evolutionary Hypothesis', *Science*, 197(4302), pp. 493–495. doi: 10.1126/science.301659.

- Valmaggia, L. R., Latif, L., Kempton, M. J. and Rus-Calafell, M. (2016) 'Virtual reality in the psychological treatment for mental health problems: An systematic review of recent evidence', *Psychiatry Research*. Elsevier, 236, pp. 189–195. doi: 10.1016/j.psychres.2016.01.015.
- Van Ombergen, A., Van Rompaey, V., Maes, L. K., Van de Heyning, P. H. and Wuyts, F. L. (2016) 'Mal de débarquement syndrome: a systematic review', *Journal of Neurology*. Springer Berlin Heidelberg, 263(5), pp. 843–854. doi: 10.1007/s00415-015-7962-6.
- Verschure, P. F. M. J. (2011) 'Neuroscience, virtual reality and neurorehabilitation: Brain repair as a validation of brain theory', *Proceedings of the Annual International Conference of the IEEE Engineering in Medicine and Biology Society, EMBS*, pp. 2254–2257. doi: 10.1109/IEMBS.2011.6090428.
- Villard, S. J., Flanagan, M. B., Albanese, G. M. and Stoffregen, T. A. (2008) 'Postural instability and motion sickness in a virtual moving room.', *Human factors*, 50(2), pp. 332–45. doi: 10.1518/001872008X250728.
- Viñas-Diz, S. and Sobrido-Prieto, M. (2016) 'Virtual reality for therapeutic purposes in stroke: A systematic review', *Neurología (English Edition)*, 31(4), pp. 255–277. doi: 10.1016/j.nrleng.2015.06.007.
- Wada, T., Fujisawa, S. and Doi, S. (2016) 'Analysis of driver's head tilt using a mathematical model of motion sickness', *International Journal of Industrial Ergonomics*, (November). doi: 10.1016/j.ergon.2016.11.003.
- Wada, T. and Yoshida, K. (2016) 'Effect of passengers' active head tilt and opening/closure of eyes on motion sickness in lateral acceleration environment of cars', *Ergonomics*. Taylor & Francis, 59(8), pp. 1050–1059. doi: 10.1080/00140139.2015.1109713.

1 Wang, J. and Lewis, R. F. (2016) 'Contribution of intravestibular sensory conflict to motion
2 sickness and dizziness in migraine disorders', *Journal of Neurophysiology*, 116(4), pp.
3 1586–1591. doi: 10.1152/jn.00345.2016.
4

5
6
7 Warwick-Evans, L. and Beaumont, S. (1995) 'An Experimental Evaluation of Sensory Conflict
8 Versus Postural Control Theories of Motion Sickness', *Ecological Psychology*, 7(3), pp.
9 163–179. doi: 10.1207/s15326969eco0703_1.
10
11

12
13
14 Weech, S. and Troje, N. F. (2017) 'Vection Latency Is Reduced by Bone-Conducted Vibration
15 and Noisy Galvanic Vestibular Stimulation', *Multisensory Research*, 30(1), pp. 65–90.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2

- 1 Wright, W. G. (2014) 'Using virtual reality to augment perception, enhance sensorimotor
2 adaptation, and change our minds.', *Frontiers in systems neuroscience*, 8(April), p. 56.
3
4 doi: 10.3389/fnsys.2014.00056.
5
6
7 Young, L. R., Dichgans, J., Murphy, R. and Brandt, T. (1973) 'Interaction Of Optokinetic And
8 Vestibular Stimuli In Motion Perception', *Acta Oto-Laryngologica*. Taylor & Francis,
9
10 76(1–6), pp. 24–31. doi: 10.3109/00016487309121479.
11
12
13 Zacharias, G. L. and Young, L. R. (1981) 'Influence of combined visual and vestibular cues on
14 human perception and control of horizontal rotation', *Experimental Brain Research*.
15
16 Springer-Verlag, 41(2), pp. 159–171. doi: 10.1007/BF00236605.
17
18
19 Zanbaka, C., Lok, B., Babu, S., Xiao, D., Ulinski, A. and Hodges, L. F. (2004) 'Effects of
20 travel technique on cognition in virtual environments', *Proceedings - Virtual Reality
21 Annual International Symposium*, pp. 149–156. doi: 10.1109/VR.2004.1310068.
22
23
24
25
26
27
28 Zu Eulenburg, P., Caspers, S., Roski, C. and Eickhoff, S. B. (2012) 'Meta-analytical definition
29 and functional connectivity of the human vestibular cortex', *NeuroImage*. Elsevier Inc.,
30
31 60(1), pp. 162–169. doi: 10.1016/j.neuroimage.2011.12.032.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1. Symptoms and Physiological Changes in Cybersickness.

Symptoms of cybersickness falling under each category of Kennedy et al.’s (1993) Simulator Sickness Questionnaire, and reported physiological changes caused by cybersickness.

Symptoms of cybersickness (according to the Simulator Sickness Questionnaire Categories)			Physiological changes in cybersickness	
<i>Nausea</i>	<i>Oculomotor</i>	<i>Disorientation</i>	<i>Increases in</i>	<i>Decreases in</i>
Discomfort Increased Salivation Sweating Nausea Difficulty Concentrating Stomach Awareness Burping	Discomfort Fatigue Headache Eyestrain Difficulty Focusing Difficulty Concentrating Blurred Vision	Difficulty Focusing Nausea Fullness of head Blurred vision Dizziness Vertigo	Heart rate Respiration rate Skin conductance Gastric activity Blinks EEG Alpha power EEG Beta power EEG Gamma power	Photoplethysmogram Skin temperature Heart period EEG Theta power

Table 2. Theories of Cybersickness

A brief overview of motion sickness theories commonly applied to cybersickness.

Theory	Authors	Key aspects
Sensory conflict theories	Reason & Brand 1975	Sensory signals which do not match stored sensory signals generate a mismatch signal, triggering motion sickness
	Oman 1988	Sensory conflicts increase a mismatch vector, triggering motion sickness
	Bles et al. 1998	Conflicts between the sensed and predicted gravitational verticals trigger motion sickness
Other theories	Riccio & Stoffregen 1991	Postural instability causes motion sickness